

On the Motion of b -quark inside B meson: the Parton Model *vs.* the ACCMM Model

C. S. Kim^{a*}, Yeong Gyun Kim^{b†} and Kang Young Lee^{a‡§}

^a *Department of Physics, Yonsei University, Seoul 120-749, Korea*

^b *Department of Physics, Korea Advanced Institute of Science and Technology, Taejon 305-701,
Korea*

(February 1, 2008)

Abstract

We study the distribution function of the b -quark for inclusive semileptonic B decays in the framework of the parton model. We compare the functional behavior of the fragmentation inspired function and the Gaussian ACCMM function, and discuss their validity. Using the HQET parameters, we can critically test these models.

Typeset using REVTeX

*kim@cskim.yonsei.ac.kr, cskim@kekvox.kek.jp

†ygkim@chep6.kaist.ac.kr

‡kylee@chep5.kaist.ac.kr

§Department of Physics, KAIST, Taejon 305-701, Korea

I. INTRODUCTION

Studies of inclusive semileptonic decays of B mesons have drawn our interest as a testing ground for the motion of b -quarks inside hadrons. We have the differential decay width for the process $B \rightarrow X_q l \nu$

$$d\Gamma \sim W^{\mu\nu} L_{\mu\nu} \frac{d^3 p_l}{(2\pi)^3 2E_l} \frac{d^3 p_\nu}{(2\pi)^3 2E_\nu}, \quad (1)$$

where $L_{\mu\nu}$ is the leptonic tensor and the hadronic tensor $W^{\mu\nu}$ is defined as

$$W^{\mu\nu} = \frac{1}{2} \sum_{spins} \int \frac{d^3 p_X}{(2\pi)^3 2E_X} (2\pi)^4 \delta^4(p_B - q - p_X) \langle B | J^{\mu\dagger} | X \rangle \langle X | J^\nu | B \rangle. \quad (2)$$

Since the B meson is heavy, the momentum transfer of the decay is larger than the typical energy scale of hadronic binding, $\sim \Lambda_{QCD}$, in most regions of phase space. This enables us to formulate the inclusive semileptonic decay process in an analogous manner to the well-known method of the deep inelastic scattering (DIS). Light-cone dynamics dominates the decay process and the hadronic form factors show the remarkable feature of scaling in the absence of the relevant corrections. So the parton picture should work well, where the modeling of the decay process is possible by summing the partonic decay processes weighted by the distribution of the b -quark momentum inside the B meson. Bareiss and Paschos [1] noticed this point and they formulated the inclusive decays along the lines of analyzing DIS in the parton picture. The general expression for the hadronic tensor in terms of invariants is given by

$$\begin{aligned} W^{\mu\nu}(q^2, q \cdot p_B) = & -g^{\mu\nu} W_1(q^2, q \cdot p_B) + p_B^\mu p_B^\nu \frac{W_2(q^2, q \cdot p_B)}{m_B^2} \\ & - i\epsilon^{\mu\nu\alpha\beta} p_{B\alpha} q_\beta \frac{W_3(q^2, q \cdot p_B)}{2m_B^2} + q^\mu q^\nu \frac{W_4(q^2, q \cdot p_B)}{m_B^2} \\ & + (p_B^\mu q^\nu + p_B^\nu q^\mu) \frac{W_5(q^2, q \cdot p_B)}{2m_B^2} - (p_B^\mu q^\nu - p_B^\nu q^\mu) \frac{W_6(q^2, q \cdot p_B)}{2m_B^2}. \end{aligned} \quad (3)$$

Following the parton picture of Feynman [2], we let the probability of finding a b -quark in a B meson carrying a fraction x of the meson momentum in the infinite momentum frame be

the distribution function $f(x)$. The decay width is then obtained by incoherently summing the partonic decay widths weighted by $f(x)$ in that frame. Thus we immediately find that

$$\begin{aligned}
W_1(q^2, q \cdot p_B) &= \frac{1}{2} (f(x_+) + f(x_-)) \\
W_2(q^2, q \cdot p_B) &= \frac{2}{x_+ - x_-} (x_+ f(x_+) - x_- f(x_-)) \\
W_3(q^2, q \cdot p_B) &= -\frac{1}{x_+ - x_-} (f(x_+) - f(x_-)) \\
W_{4,5,6}(q^2, q \cdot p_B) &= 0
\end{aligned} \tag{4}$$

after neglecting the lepton masses where

$$x_{\pm} = \frac{q_0 \pm \sqrt{\mathbf{q}^2 + m_q^2}}{m_B} \tag{5}$$

where m_q is the final state quark mass in $b \rightarrow qe\nu$. This approach has been reformulated on the light-cone in Ref. [3–5] and holds in any frame. The QCD radiative correction of the parton model was studied in Ref. [5]. The idea of the parton model was also applied to the exclusive semileptonic decays of B meson in Ref. [6].

Recently the heavy quark effective theory (HQET) has been applied to inclusive semileptonic decays of B mesons and brought great progress [7–11]. With the expansion for the decay rate in inverse power of b -quark mass, a systematic study is possible and one obtains more information about the hadronic matrix elements than before. Perhaps one of the most important results of the HQET analysis for the inclusive decays is the absence of the contribution of the $1/m_b$ correction terms. With the help of the $1/m_b$ expansion, we can relate the moments of the charged lepton energy spectrum to the matrix elements for the operator expansion of the current product.

The $1/m_b$ structure of the inclusive parton model has recently been analyzed in Ref. [12]. When the parton model approach motivated by DIS is applied to the analysis of B decays, the modeling boils down to introducing the distribution function of Eq. (4), which describe the b -quark's motion inside the B meson. In other words, in the study of the inclusive semileptonic B decays, the problem of which model we choose corresponds to the problem of choice of the distribution function. In general two models are known which predict the

lepton energy spectrum of inclusive decays of B meson. Altarelli et al. (ACCMM) [13] treat the b -quark and the spectator quark as quasi-free particles with a Gaussian spectrum of Fermi momentum in the B rest frame. Bareiss and Paschos [1,14] notice that the b -quark distribution inside a B meson in the infinite momentum frame (IMF) is related to the fragmentation of a b -quark into a B meson by crossing symmetry of the spectator, and so their functional behaviors are similar to each other. Thus they chose Peterson's fragmentation function [15] as the distribution function of b -quark, which is the default choice in Lund Monte Carlo programs. Both models get results at the hadronic level after convolution of the quark level spectrum. The difference between the two models mainly comes from the weight function which is corresponding to the distribution function $f(x)$ of b -quark momentum in the parton model approach.

In this paper, we study both models from the viewpoint of the parton model picture to compare their distribution functions with each other. For explicit comparison of the distribution functions of both models, we derive the distribution functions described by the same definition in the same reference frame. Moreover, with the help of the $1/m_b$ study suggested in Ref. [12] we attempt to test the models by extracting the HQET parameters, the effective mass of light degrees of freedom $\bar{\Lambda}$ and the average kinetic energy of b -quark μ_π^2 . We show that known bounds on $\bar{\Lambda}$ and μ_π^2 constrain the model parameters and may even rule out the model itself. In section 2, we review the parton model approach first. In this approach Peterson's function and its improved form by Lee and Kim [12] is taken as the distribution function. In the ACCMM model, the distribution of the b -quark is determined by the Gaussian distribution for the Fermi motion of the spectator quark in the B meson rest frame. In section 3, we derive the distribution function of the ACCMM model in the infinite momentum frame from the Gaussian function, which allows a consistent comparison of the two models. The behavior of the distribution functions is investigated and the heavy quark effective theory (HQET) parameters are obtained in terms of the parameters of each model. Our discussions and conclusions are summarized in section 4.

II. THE PARTON MODEL

We first review the parton model approach applied to analyze inclusive semileptonic B meson decays. As described in the previous section, the distribution function $f(x)$ is defined as the probability of finding a b -quark in a B meson carrying a fraction x of the meson momentum in the infinite momentum frame. Thus the decay probability is determined by the momentum of the b -quark and has the value of $f(x)dx$. And we write the Lorentz invariant decay width as follows

$$E_B d\Gamma(B \rightarrow X_q e \nu) = \int dx f(x) E_b d\Gamma(b \rightarrow q e \nu), \quad (6)$$

with the relation $p_b = xp_B$. Since B mesons are heavy, light-cone dynamics dictates the inclusive semileptonic decay analogous to that of DIS. The universal distribution function of QCD is interpreted as the structure function of the b -quark in the parton model, which determines the distribution of the light-cone projection of the b -quark momentum inside the B meson. We directly derive the triple differential decay rate as

$$\frac{d\Gamma}{dE_l dq^2 dq_0} = \frac{G_F^2 |V_{qb}|^2}{4\pi^3} \frac{q_0 - E_l}{\sqrt{\mathbf{q}^2 + m_q^2}} [x_+ f(x_+) (2E_l - m_B x_-) + (x_+ \leftrightarrow x_-)] \quad (7)$$

where $x_{\pm}$ is defined in the previous section. Jin and Paschos [3] argued that the contribution of $f(x_-)$ is expected to be relatively small in the kinematic region we are interested in, and can be neglected. This is a consequence of the fact that the light-cone distribution function is sharply peaked around $x \sim m_b/m_B$.

For a more quantitative analysis, we have to introduce a method analogous to the HQET. In this parton model approach, information on the nonperturbative dynamics inside the B meson is encoded in the distribution function. Therefore, it is crucial to determine precisely the form of the distribution function in this approach. Unfortunately we don't presently know how to calculate the true form of it. However, we have some knowledge about the line shape function of QCD from the help of HQET and the operator product expansion (OPE). Since the mass of the b -quark does not appear in this parton model framework, we

have to define it within this framework in order to express the model in terms of the inverse power expansion of the b -quark mass. With properly defined m_b , the parton model is well represented by an $1/m_b$ expansion [12]. We define

$$m_b \equiv \langle x \rangle m_B, \quad (8)$$

where $\langle x \rangle = \int dx \, x f(x)$. With this definition as the starting point, we can relate the parton model to HQET and compute the effective mass of the light degrees of freedom in a B meson, $\bar{\Lambda}$.

In the HQET framework, the triple differential decay rate for $B \rightarrow X_q l \nu$ shows scaling behavior [8–10]:

$$\frac{d\Gamma}{dE_l dq^2 dq_0} = \frac{G_F^2 |V_{qb}|^2}{192\pi^3} \frac{2}{\bar{\Lambda}} F(x_B) \frac{24(q_0 - E_l)(2m_b E_l - q^2)}{m_b - q_0}, \quad (9)$$

where

$$x_B = -\frac{m_b^2 + q^2 - 2m_b q_0}{2\bar{\Lambda}(m_b - q_0)}. \quad (10)$$

Apart from kinematic factors, the triple differential decay rate is completely determined by one scaling variable x_B . The scaling behavior of QCD will be violated by perturbative and nonperturbative (higher twist) corrections. If we take the final state quark to be u -quark and so $m_q = 0$ hereafter, the scaling form is derived from Eq. (6)

$$\frac{d\Gamma}{dE_l dq^2 dq_0} \approx \frac{G_F^2 |V_{ub}|^2}{192\pi^3} \frac{2}{m_B} f(x_+) \frac{24(q_0 - E_l)(2m_b E_l - q^2)}{m_b - q_0} \quad (11)$$

where m_b is as defined in Eq. (7). From Eqs. (8) and (10), the line shape function $F(x_B)$ of QCD is related to the distribution function $f(x_+)$ of the parton model after appropriate variable changes

$$\frac{1}{m_B} f(x_+) = \frac{1}{\bar{\Lambda}} F(x_B). \quad (12)$$

Next we intend to study their functional behavior. Following the notation of Bigi et al. [9,10], a general form for the singular expansion of the distribution function is written as

$$F(x_B) = \sum_{n=0}^{\infty} \frac{(-1)^n}{n!} a_n \delta^{(n)}(x_B), \quad (13)$$

where moments of the equation:

$$a_n = \int dx_B x_B^n F(x_B), \quad n = 0, 1, \dots \quad (14)$$

Using the HQET and the OPE, we parametrize the moments with the forward scattering matrix elements of B meson. Especially we know that $a_0 = 1$ from b -number conservation, and that $a_1 = 0$ up to $\mathcal{O}(1/m_b^2)$, which indicates the lack of $\mathcal{O}(1/m_b)$ corrections. The value of the first moment $a_1 = 0$ is related to the value of the b -quark mass m_b or the parameter $\bar{\Lambda}$.

The second moment of the spectrum is related to the average kinetic energy of b -quark inside the B meson and usually represented by the parameter μ_π^2 , defined by

$$a_2 = \frac{1}{3\bar{\Lambda}^2} \frac{1}{2m_B} \langle B | \bar{b} \pi^2 b | B \rangle \equiv \frac{\mu_\pi^2}{3\bar{\Lambda}^2}, \quad (15)$$

where $\pi_\alpha = iD_\alpha - m_b v_\alpha$ with D_α the covariant derivative and v_α the four-velocity of the B meson. After the variable changes of Eq. (11), we can calculate the moments for the distribution function in the framework of the parton model. Under the definition of m_b in Eq. (7), we have $a_1 = 0$. The second moment is derived as

$$a_2 = \int_{-\infty}^1 dx_B x_B^2 F(x_B) = \left(\frac{m_B}{\bar{\Lambda}} \right)^2 \int_0^1 dx \left(x - \frac{m_b}{m_B} \right)^2 f(x) = \left(\frac{m_B}{\bar{\Lambda}} \right)^2 (\langle x^2 \rangle - \langle x \rangle^2). \quad (16)$$

We first used Peterson's fragmentation function as the distribution function of b -quark following the original work of Bareiss and Paschos [1], and obtained the value for $a_2 \sim 0.76$, with the model parameter $\epsilon = 0.004$ and the mass of B meson $m_B = 5.3$ GeV, which agrees well with the QCD sum rule result of Ref. [16]. As pointed out in the Ref. [12], however, this function gives rather larger $\bar{\Lambda}$ and μ_π^2 , and we need to find a better functional form or some modification for Peterson's function. Jin and Paschos [3] also pointed out that the distribution function should be more sharply peaked than Peterson's function. By improving Peterson's arguments, Lee and Kim [12] suggested a modified form of Peterson's function

$$f_Q^{\text{new}}(x) = \frac{N_Q}{x(1 + \alpha_Q - \frac{1}{x} - \frac{\epsilon_Q}{1-x})^2}. \quad (17)$$

Using this new form, we calculate $\bar{\Lambda}$ and μ_π^2 by varying the input parameters α_Q and ϵ_Q .

Fig. 1 shows the behavior of the distribution function with varying α_Q and ϵ_Q . The larger the value of α_Q , the sharper the functional behavior. And the smaller the value of ϵ_Q , the sharper the function and the closer the location of peak to 1. We also show the parameter space of (α_Q, ϵ_Q) for allowed $\bar{\Lambda}$ and μ_π^2 in Fig. 2. For the values of μ_π^2 , there is some theoretical controversy which we will discuss later in section 4. We take the range of $\bar{\Lambda}$ and μ_π^2 obtained by the HQET and QCD sum rules in Refs. [16,17]

$$\bar{\Lambda} = 0.4 \sim 0.6,$$

$$\mu_\pi^2 = 0.6 \pm 0.1 \quad [16], \quad \text{and} \quad \mu_\pi^2 = 0.10 \pm 0.05 \quad [17]. \quad (18)$$

Ball and Braun [16] calculated μ_π^2 using the QCD sum rule approach and obtained $\mu_\pi^2 = 0.60 \pm 0.10 \text{ GeV}^2$ for B -meson, while Neubert [17] obtained $-\lambda_1 = 0.10 \pm 0.05 \text{ GeV}^2$. We find that there exists a region in the (α_Q, ϵ_Q) plane that gives values of $\bar{\Lambda}$ and μ_π^2 in agreement with those of Ball and Braun [16]. If Neubert's prediction [17] for the value of μ_π^2 is correct, there is no common region in which the model parameters give the values of $\bar{\Lambda}$ and μ_π^2 in Eq. (17), and we may conclude that this functional form is not appropriate for inclusive semileptonic B decays.

III. THE ACCMM MODEL

The ACCMM model [13] considers the B meson as consisting of a b -quark and a spectator quark. The spectator quark is treated as a quasi-free particle having definite mass m_{sp} , and momentum $\mathbf{p}$, while the b -quark is treated as a virtual particle of invariant mass m_b

$$m_b^2(p) = m_B^2 + m_{sp}^2 - 2m_B \sqrt{p^2 + m_{sp}^2}, \quad (19)$$

where m_B is the B meson mass and $p \equiv |\mathbf{p}|$ in the B meson rest frame. The b -quark energy is given by

$$\begin{aligned}
E_b(p) &= \sqrt{p^2 + m_b^2(p)} \\
&= m_B - \sqrt{p^2 + m_{sp}^2} .
\end{aligned}
\tag{20}$$

The distribution of the spectator momentum is usually assumed to be a Gaussian with an adjustable parameter p_F

$$\Phi(p) = \frac{4}{\sqrt{\pi} p_F^3} \exp\left(-\frac{p^2}{p_F^2}\right)
\tag{21}$$

with the normalization

$$\int_0^\infty dp \, p^2 \Phi(p) = 1.
\tag{22}$$

After convoluting the distribution of the spectator momentum with the subprocess of the virtual b -quark decay in a frame where its momentum is $-\mathbf{p}$, we can obtain the lepton energy spectrum at the hadronic level. This model has been extensively used in the analysis of the charged lepton energy spectrum in semileptonic decays and numerically reproduce the inclusive spectra well. In the light of QCD, several authors analyzed the model [18] and suggested a way to reproduce the pattern of the QCD description of the heavy quark motion. In this paper we derive the distribution function $f(x)$ in the IMF from the given Gaussian distribution in the B rest frame, $\Phi(p)$, in order to study the ACCMM model by comparing with the parton model approach.

Let's choose a coordinate system where the B meson is at the origin in B rest frame. We boost the B meson along the positive z direction with the velocity β . As is well known, for the B meson at rest we have the boosted energy of B meson as

$$E_B^* = \gamma E_B = \gamma m_B,
\tag{23}$$

where $\gamma = 1/\sqrt{1 - \beta^2}$, and that of b -quark as

$$E_b^*(p) = \gamma(E_b(p) - \beta p_z),
\tag{24}$$

where p is the magnitude of the spatial momentum of the b -quark and p_z is the z component of $\mathbf{p}$ in the B -rest frame. Now we calculate the energy ratio between b -quark and B meson in this moving frame

$$\frac{E_b^*}{E_B^*} = \frac{\gamma(E_b - \beta p_z)}{\gamma m_B}. \quad (25)$$

In the limit of $\beta \rightarrow 1$, $\gamma \rightarrow \infty$, we define the ratio as

$$\frac{E_b^*}{E_B^*} = \frac{(E_b - p_z)}{m_B} \equiv x, \quad (26)$$

which represents the ratio of b -quark energy to B meson energy in the IMF.

Because no specific direction for $\mathbf{p}$ is preferred in B meson decay, the value of p_z is distributed from $-p$ to $+p$ with equal probability for a fixed value of p . Then the value of x can be from $x_{min}(p)$ to $x_{max}(p)$ with equal weight for a fixed p , where

$$\begin{aligned} x_{min}(p) &= \frac{1}{m_B}(E_b(p) - p), \\ x_{max}(p) &= \frac{1}{m_B}(E_b(p) + p), \end{aligned} \quad (27)$$

with the weight $m_B/2p$, because $x_{max}(p) - x_{min}(p) = 2p/m_B$. Each value of p in the B meson rest frame corresponds to a distribution of x in the IMF such that the probability of finding a b -quark in a B meson carrying a fraction x of the meson momentum in the IMF is obtained by intergrating over p with Gaussian weight, $\Phi(p)$,

$$\begin{aligned} f(x) &= \int_0^{p_{max}} dp p^2 \Phi(p) \cdot \frac{m_B}{2p} \theta(x - x_{min}(p)) \theta(x_{max}(p) - x) \\ &= \int_{p_{min}}^{p_{max}} dp p^2 \Phi(p) \cdot \frac{m_B}{2p} \end{aligned} \quad (28)$$

where

$$p_{min} = \frac{|m_B^2(1-x)^2 - m_{sp}^2|}{2m_B(1-x)} \quad (29)$$

is obtained by the conditions $x_{min}(p) < x < x_{max}(p)$. We take the value of p_{max} to be infinity as far as the integration is concerned, because the area of phase space for $p > p_{max}$ is negligible. After the integration over p , we obtain

$$f(x) = \frac{m_B}{\sqrt{\pi} p_F} \exp \left(-\frac{1}{4} \left(\frac{\rho p_F}{m_B(1-x)} - \frac{m_B}{p_F}(1-x) \right)^2 \right), \quad (30)$$

where $\rho = m_{sp}^2/p_F^2$. Using the Eq. (12) and the relation, $(1 - x_+) = \frac{\bar{\Lambda}}{m_B}(1 - x_B)$, we find Eq. (30) becomes Eq. (30) of Ref. [9], the light-cone distribution function for the ACCMM

model derived by Bigi *et al.* We show the functional behavior of $f(x)$ in Fig. 3 by varying the parameters p_F and m_{sp} . Note that the shape of the function $f(x)$ depends on two parameters m_{sp} and p_F . The peak position is determined by m_{sp} :

$$x_{peak} = \frac{m_B - m_{sp}}{m_B}, \quad (31)$$

while the Fermi motion parameter p_F is related to the width of the distribution $f(x)$, i.e. the larger the value of p_F , the broader the distribution $f(x)$. One can easily check the normalization of $f(x)$

$$\int_0^1 f(x) dx = \int_0^{p_{max}} dp p^2 \Phi(p) = 1. \quad (32)$$

The first moment of $f(x)$ is directly related to the average b -quark energy in the B rest frame,

$$\langle x \rangle \equiv \int_0^1 dx x f(x) = \int_0^{p_{max}} dp p^2 \Phi(p) \frac{E_b(p)}{m_B} \equiv \frac{\langle E_b(p) \rangle_{B-rest}}{m_B}, \quad (33)$$

because $m_b \equiv \langle x \rangle m_B$ in the parton model approach we get the relation $m_b \equiv \langle E_b \rangle_{B-rest}$, which reminds us of Csáki and Randall's definition of the b -quark mass¹ for the ACCMM model with respect to the HQET.

We thus have derived a distribution function $f(x)$ of Eq. (28) for the parton model approach from a Gaussian distribution in the B rest frame. Following the method discussed in section 2, we can calculate the moments of the line shape function and the parameters $\bar{\Lambda}$ and μ_π^2 for given model parameters. First we calculate $\bar{\Lambda}$:

$$\begin{aligned} \bar{\Lambda} &\equiv m_B - m_b = m_B - \langle E_b \rangle_{B-rest} \\ &= m_B - \int_0^\infty dp p^2 \Phi(p) (m_B - \sqrt{p^2 + m_{sp}^2}) \\ &= \langle \sqrt{p^2 + m_{sp}^2} \rangle_{B-rest}. \end{aligned} \quad (34)$$

¹ Csaki and Randall [18] showed that the predictions of the ACCMM model agree well with those of the heavy quark effective theory with a definition of the b -quark mass, $m_b \equiv \langle E_b(p) \rangle_{B-rest}$.

From Eq. (14), μ_π^2 is given as

$$\mu_\pi^2 = 3\bar{\Lambda}^2 a_2 = 3m_B^2 \left(\langle x^2 \rangle - \frac{m_b^2}{m_B^2} \right). \quad (35)$$

And using

$$\begin{aligned} \langle x^2 \rangle &= \int_0^1 dx \, x^2 f(x) \\ &= \frac{1}{m_B^2} (\langle E_b^2 \rangle_{B-rest} + \frac{1}{3} \langle p^2 \rangle_{B-rest}), \end{aligned} \quad (36)$$

we get

$$\mu_\pi^2 = 3(\langle E_b^2 \rangle_{B-rest} - \langle E_b \rangle_{B-rest}^2) + \langle p^2 \rangle_{B-rest}. \quad (37)$$

Finally with the following relations,

$$\langle E_b^2 \rangle_{B-rest} - \langle E_b \rangle_{B-rest}^2 = \langle p^2 \rangle_{B-rest} + m_{sp}^2 - \langle \sqrt{p^2 + m_{sp}^2} \rangle_{B-rest}^2 \quad (38)$$

and

$$\langle p^2 \rangle_{B-rest} = \int_0^\infty dp \, p^2 \Phi(p) \, p^2 = \frac{3}{2} p_F^2, \quad (39)$$

we obtain

$$\mu_\pi^2 = 6p_F^2 + 3m_{sp}^2 - 3\langle \sqrt{p^2 + m_{sp}^2} \rangle_{B-rest}^2. \quad (40)$$

We show the parameter space of (p_F, m_{sp}) for given $\bar{\Lambda}$ and μ_π^2 in Fig. 4. Values of (p_F, m_{sp}) in the shaded region A satisfy the condition of Eq. (17) given in the Ref. [16], while values in the region B satisfy the condition of Eq. (17) given in the Ref. [17]. We can determine the model parameters p_F and m_{sp} , once we know the correct value of the HQET parameter μ_π^2 .

IV. DISCUSSION AND CONCLUSIONS

We have studied two phenomenological models (QCD parton model and ACCMM model) for the analysis of inclusive semileptonic B decays in the context of the parton model approach. In this framework, the core of modeling is the choice of a distribution function of

the b -quark inside a B meson. The light-cone dominance enables us to describe the triple differential decay rate in terms of a single function, the light-cone distribution function up to kinematic factors, as the form given in Eqs. (9) and (11). The distribution function in ACCMM model comes from the Gaussian distribution of the b -quark momentum at the B rest frame. Bigi *et al.* [9] obtained the light-cone distribution function for the ACCMM model by reading off it from the photon energy spectrum of $B \rightarrow X_s \gamma$ decays. We calculate it here by counting the probability of finding a b -quark with the boosted energy ratio x between b -quark and B meson into the infinite momentum frame within the framework of the parton model. We find that our result of Eq. (30) exactly reproduce the light-cone distribution function given by Bigi *et al.*, Eq. (30) of Ref. [9]. This agreement is originated from the fact that the ACCMM model treats the b -quark as a quasi-free particle at the decay rate level by classical treatment as we do not consider the perturbative QCD corrections in this paper.

Even though we cannot calculate the distribution function from QCD completely, some model independent information can be obtained with the help the HQET. When we expand this function with infinite number of moments, we cannot know all the moments while the HQET give some informations for a few moments. We note that since this is an singular expansion, we cannot construct a well-behaved function with only a few moments. In spite of that, the information is very useful for giving some constraints on phenomenological models. Investigating the distribution function, two moments of the function, corresponding to the mean value and the width, can be related to two QCD parameters, $\bar{\Lambda}$ (or m_b) and μ_π^2 . The mean value of the function strongly depends on $\bar{\Lambda}$ and very weakly on the μ_π^2 , while the width on the μ_π^2 . Actually the fact that $a_1 = 0$ define the quark mass, equivalent to $\bar{\Lambda}$ and the next moment a_2 is directly related to the μ_π^2 . With these constraints, we obtain the possible regions of model parameters for each model, as shown in Figs. 2 and 4.

Recently, it has been an important subject to obtain an accurate value of the kinetic energy, μ_π^2 ($\equiv -\lambda_1$), of the heavy quark inside B meson in connection with the heavy quark effective theory (HQET). Neubert derived [19] the field-theory version of the virial theorem

within the HQET framework and obtained the result $\mu_\pi^2 \sim 0.10 \pm 0.05 \text{ GeV}^2$, which is much smaller than the QCD sum rule calculations of Ball and Braun [16] and rather comparable to the earlier QCD sum rule result using a less sophisticated approach in Ref. [20]. However, it should be noted that Refs. [16] and [17] differ in the choice of the 3-point correlation functions used to estimate the matrix elements of interest. The difference in the numerical values obtained in these two calculations is understood in terms of the contributions of excited states, which in principle must be subtracted in any QCD sum rule analysis. In practice, this subtraction can only be done approximately. Therefore, the numerical differences between Refs. [16] and [17] indicate the limited accuracy of the QCD sum rule approach. Bigi *et al.* [21] derived an inequality between the expectation value of the kinetic energy of the heavy quark inside the hadron and that of the chromomagnetic operator, $\langle \mathbf{p}^2 \rangle \geq \frac{3}{4}(M_V^2 - M_P^2)$, which gives $\mu_\pi^2 \geq 0.36 \text{ GeV}^2$ for B meson system. However, Kapustin *et al.* [22] showed later that this lower bound could be significantly weakened by higher order perturbative corrections. By means of a QCD relativistic potential model, the value of μ_π^2 was found to be quite large as $0.44 \sim 0.46 \text{ GeV}^2$ in Ref. [23]. Besides the above theoretical calculations of μ_π^2 , Gremm *et al.* [24] extracted the average kinetic energy by comparing the prediction of the HQET with the shape of the inclusive $B \rightarrow X l \nu$ lepton energy spectrum [25] for $E_l \geq 1.5 \text{ GeV}$, in order to avoid the contamination from the secondary leptons of cascade decays of $b \rightarrow c \rightarrow s l \nu$. They obtained $-\lambda_1 = 0.19 \pm 0.10 \text{ GeV}^2$. Combining the experimental data on the inclusive decays of $D \rightarrow X e \nu$, $B \rightarrow X e \nu$ and $B \rightarrow X \tau \nu$, Ligeti *et al.* [26] derived the bound of $\mu_\pi^2 \leq 0.63 \text{ GeV}^2$ if $\bar{\Lambda} \geq 0.240 \text{ GeV}$, or $\mu_\pi^2 \leq 0.10 \text{ GeV}^2$ if $\bar{\Lambda} \geq 0.500 \text{ MeV}$. Li *et al.* [27] obtained the value of $-\lambda_1$ centered at 0.71 GeV^2 from the analysis of the inclusive radiative decay $B \rightarrow X_s \gamma$ [28] within the perturbative QCD framework. Related with the comparison of various theoretical calculations of μ_π^2 , we note that Ref. [29] emphasizes that one has to be careful when comparing the values of $-\lambda_1$ obtained using different theoretical methods. The mixing of the operator for the heavy quark kinetic energy with the identity operator occurs at the two-loop order and the parameter λ_1 requires a non-perturbative

subtraction.

Note also that the functional behavior of Refs. [12] and [13] is different from each other while both functions satisfy the constraints. Even though two moments, a_1 , a_2 , which correspond to the mean values and the widths of the distribution functions, agree with each other, their full behaviors do not necessarily agree. For the two models studied in this paper, we find that the modified Peterson's function in Fig. 1 has a rather longer tail than the distribution function derived from the Gaussian function of the Fermi momentum defined in the ACCMM model in Fig. 3. We cannot say which model is better, at least at this stage. Both models can give proper values for the b -quark mass and the width of the charged lepton energy spectrum, both theoretically and phenomenologically. For further tests, we must fit them to the experimental data more precisely after more data is accumulated. For this purpose the photon energy spectrum of the inclusive radiative decay $B \rightarrow X_s \gamma$ would be more useful. The photon energy spectrum of this process is known to be more sensitive to the behavior of the distribution function than the charged lepton energy spectrum of the inclusive semileptonic decays [12,27], since the charged lepton energy spectrum is contaminated by kinematics.

ACKNOWLEDGMENTS

We would like to thank M. Drees for useful suggestions and helpful comments. The work of CSK was supported in part by Non-Directed-Research-Fund, KRF in 1997, in part by the CTP, Seoul National University, in part by Yonsei University Faculty Research Fund of 1997, in part by the BSRI Program, Ministry of Education (MOE), Project No. BSRI-97-2425, and in part by the KOSEF-DFG large collaboration project, Project No. 96-0702-01-01-2. YGK was supported by CTP of SNU. KYL is a postdoctoral fellow supported by Korea Research Foundation (KRF) and Research University Fund of College of Science at Yonsei University by MOE of Korea.

REFERENCES

- [1] A. Bareiss and E. A. Paschos, Nucl. Phys. **B 327**, 353 (1989).
- [2] R. P. Feynman, Phys. Rev. Lett. **23**, 1415 (1969).
- [3] C. H. Jin and E. A. Paschos, preprint DO - TH 95/07 (1995).
- [4] C. H. Jin, Phys. Rev. **D 56**, 2928 (1997).
- [5] C. H. Jin and E. A. Paschos, hep-ph/9704405 (1997).
- [6] C. S. Kim, J. K. Kim, Y. G. Kim and K. Y. Lee, Int. J. Mod. Phys. **A 12**, 2031 (1997).
- [7] J. Chay, H. Georgi and B. Grinstein, Phys. Lett. **B 247**, 399 (1990); I.I. Bigi, M.A. Shifman, N.G. Uraltsev and A.I. Vainshtein, Phys. Rev. Lett. **71**, 496 (1993); T. Mannel, Nucl. Phys. **B 413**, 396 (1994).
- [8] A. V. Manohar and M. B. Wise, Phys. Rev. **D 49**, 1310 (1994); B. Blok, L. Koyrakh, M. Shifman and A. I. Vainstein, Phys. Rev. **D 49**, 3356 (1994).
- [9] I. I. Bigi, M. A. Shifman, N. G. Uraltsev and A. L. Vainstein, Phys. Lett. **B 328**, 431 (1994).
- [10] I. I. Bigi, M. A. Shifman, N. G. Uraltsev and A. L. Vainstein, Int. J. Mod. Phys. **A 9**, 2467 (1994).
- [11] M. Neubert, Phys. Rev. **D 49**, 3392 (1994); *ibid* 4623.
- [12] K. Y. Lee and J. K. Kim, Phys. Lett. **B 377**, 153 (1996).
- [13] G. Altarelli, N. Cabbibo, G. Corbo, L. Maiani and G. Martinelli, Nucl. Phys. **B 208**, 365 (1982).
- [14] C. H. Jin, W. F. Palmer and E. A. Paschos, Phys. Lett. **B 329**, 364 (1994); C. H. Jin, W. F. Palmer and E. A. Paschos, preprint DO-TH 93/31 (1993).
- [15] C. Peterson, D. Schlatter, I. Schmitt and P. M. Zerwas, Phys. Rev. **D 27**, 105 (1983).

- [16] P. Ball and V. M. Braun, Phys. Rev. **D 49**, 2472 (1994).
- [17] M. Neubert, Phys. Lett. **B 389**, 727 (1996).
- [18] G. Baillie, Phys. Lett. **B 324**, 446 (1994); C. Csáki and L. Randall, Phys. Lett. **B 324**, 451 (1994).
- [19] M. Neubert, Phys. Lett. **B 322**, 419 (1994).
- [20] V. Eletsky and E. Shuryak, Phys. Lett. **B 276**, 191 (1992).
- [21] I.I. Bigi, M.A. Shifman, N.G. Uraltsev and A.I. Vainshtein, Phys. Rev. **D 52**, 196 (1995).
- [22] A. Kapustin, Z. Ligeti, M.B. Wise and B. Grinstein, Phys. Lett. **B 375**, 327 (1996).
- [23] D.S. Hwang, C.S. Kim and W. Namgung, Z. Phys. **C 69**, 107 (1995); Phys. Rev. **D 53**, 4951 (1996); Phys. Rev. **D 54**, 5620 (1996); Phys. Lett. **B 406**, 117 (1997); F. de Fazio, Mod. Phys. Lett. **A 11**, 2693 (1996).
- [24] M. Gremm, A. Kapustin, Z. Ligeti and M.B. Wise, Phys. Rev. Lett. **77**, 20 (1996).
- [25] CLEO Collaboration, J. Bartelt *et al.*, Report No. CLEO/CONF 93-19 (1993).
- [26] Z. Ligeti and Y. Nir, Phys. Rev. **D 49**, 4331 (1994).
- [27] H. Li and H.-L. Yu, Phys. Rev. **D 55**, 2833 (1997).
- [28] CLEO Collaboration, M. S. Alam *et al.*, Phys. Rev. Lett. **74**, 2885 (1995).
- [29] M. Neubert, Phys. Lett. **B 393**, 110 (1997).

FIGURES

FIG. 1. (a) Functional behavior of the improved Peterson's function with varying $\alpha_Q = 0.05, 0.085, 0.1$ for fixed $\epsilon_Q = 0.004$; (b) the same function with varying $\epsilon_Q = 0.002, 0.004, 0.006$ for fixed $\alpha_Q = 0.085$.

FIG. 2. The model parameter space in the (α_Q, ϵ_Q) plane for the QCD sum rule results for $\bar{\Lambda}$ and μ_π^2 . Values of (α_Q, ϵ_Q) in the shaded region satisfy the condition $0.4 < \bar{\Lambda} < 0.6$ and $0.5 < \mu_\pi^2 < 0.7$.

FIG. 3. (a) Functional behavior of the distribution function in the infinite momentum frame derived from the Gaussian distribution of spectator momenta with varying $p_F = 0.3, 0.5, 0.7$ GeV for fixed $m_{sp} = 0.1$ GeV; (b) the same function with varying $m_{sp} = 0.0, 0.2, 0.4$ GeV for fixed $p_F = 0.5$ GeV

FIG. 4. The model parameter space in the (p_F, m_{sp}) plane for the QCD sum rule results for $\bar{\Lambda}$ and μ_π^2 . Values of (p_F, m_{sp}) in the shaded region A satisfy the condition $0.4 < \bar{\Lambda} < 0.6$ and $0.5 < \mu_\pi^2 < 0.7$. Values of (p_F, m_{sp}) in the shaded region B satisfy the condition $0.4 < \bar{\Lambda} < 0.6$ and $0.05 < \mu_\pi^2 < 0.15$.




Fig. 1 (a)




Fig. 1 (b)




Fig. 2




Fig. 3 (a)




Fig. 3 (b)




Fig. 4